


## Rabbit antibody to VDP

<b>Code</b>	OSV00035G
<b>ID Tag</b>	Rb699-060908-G
<b>Unit size</b>	500 Å, Åµg
<b>Immunogen</b>	A synthetic peptide from human VDP conjugated to blue carrier protein was used as the antigen. The antigen is homologous in rat and mouse.
<b>Conjugate</b>	Unconjugated antibody
<b>Also known</b>	Protein USO1 homolog, Transcytosis-associated protein, TAP, General vesicular transport factor p115, USO1, VDP
<b>Host</b>	NZ white rabbit
<b>Purity</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Polyclonal, IgG
<b>Applications</b>	IHC, WB. A concentration of 10-50 µg/ml is recommended. The optimal concentration should be determined by the end user. Not yet tested in other applications.
<b>Specificity</b>	Specific for VDP.
<b>Spcs X-react.</b>	Human, Mouse, rat. Other species not yet tested.
<b>Format</b>	Lyophilised
<b>Reconstitution</b>	Reconstitute in 500 µl of sterile water. Centrifuge to remove any insoluble material.
<b>Storage</b>	Maintain the lyophilised/reconstituted antibodies frozen at -20°C for long term storage and refrigerated at 2-8°C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles.
<b>Expiry Date</b>	12 months after reconstitution
<b>Shipping</b>	This item will be shipped to you at ambient temperature in a lyophilised form.
<b>Limitation</b>	For research use only



Human Melanoma cell line C 32 was cultured overnight on round cover slides placed in a 24 well tissue culture plate. Culture media removed and washed twice with PBS before fixing with 2% formalin for 10 minutes. Cells were then washed three times with PBS and incubated with Tris 0.01M containing Triton X 0.005% for 15 minutes. Cells were washed and incubated with 100 µl of Rabbit antibody to internal region of Vesicle docking protein (VDP, USO1, TAP, Transcytosis-associated protein): IgG (OSV00035G) diluted 1:100 in the blocking buffer for 30 minutes. Wells were then washed 7 times with PBS and incubated with 100 µl of anti Rb-FITC conjugate diluted 1:100 in the blocking buffer for further 30 minutes. Cells were washed as before and nuclear counter stained with Hoechst and mounted on to slides.

---

### Related Products

OSV00018W Rabbit antibody to VDP