

Goat antibody to extracellular ASIC3 (50-100)

| | |
|-----------------------|---|
| Code | OSA00114W |
| ID Tag | Gt225-131209-WS |
| Unit size | 100 ul |
| Immunogen | A synthetic peptide from aa region 50-100 of rat ASIC3 conjugated to blue carrier protein was used as the antigen. The peptide is homologous in mouse. |
| Conjugate | Unconjugated antibody |
| Also known | Amiloride-sensitive cation channel 3, Acid-sensing ion channel 3, ASIC3, ASIC3, Testis sodium channel 1, TNaC1, TNaC1, SLNAC1, ACCN3, TNAC1 |
| Host | Goat |
| Purity | Whole serum |
| Clonality | Polyclonal |
| Isotype | Polyclonal, whole serum |
| Applications | IHC, WB. A dilution of 1 : 300 to 1 : 2000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications. |
| Specificity | Specific for ASIC3. |
| Spcs X-react. | Rat, mouse. Other species not yet tested. |
| Format | Lyophilised |
| Reconstitution | Reconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material. |
| Storage | Maintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and refrigerated at 2-8C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles. |
| Expiry Date | 12 months after reconstitution |
| Shipping | This item will be shipped to you at ambient temperature in a lyophilised form. |
| Limitation | For research use only |