

Rabbit antibody to KCNH2 (1120-1162)

ImmunogenA synthetic peptide from aa region 1120-1162 of mouse KCNH2 conjugated to blue carrier protein was used as the antigen. The peptide is homologous in rat and human.ConjugateUnconjugated antibodyAlso knownPotassium voltage-gated channel subfamily H member 2, Eag homolog, Ether-a-go-go-related gene potassium channel 1, ERG-1, Eag-related protein 1, Ether-a-go-go-related protein 1, Voltage-gated potassium channel subunit Kv11.1, ERG, ERG1, HERGHostNZ white rabbitPurityWhole serumClonalityPolyclonal
Also knownPotassium voltage-gated channel subfamily H member 2, Eag homolog, Ether-a-go-go-related gene potassium channel 1, ERG-1, Eag-related protein 1, Ether-a-go-go-related protein 1, Voltage-gated potassium channel subunit Kv11.1, ERG, ERG1, HERGHostNZ white rabbitPurityWhole serum
gene potassium channel 1, ERG-1, Eag-related protein 1, Ether-a-go-go-related protein 1, Voltage-gated potassium channel subunit Kv11.1, ERG, ERG1, HERGHostNZ white rabbitPurityWhole serum
Purity Whole serum
Clonality Polyclonal
Cionality i Dividinali
Isotype Polyclonal, whole serum
Applications IHC, WB. A dilution of 1 : 1000 to 1 : 2000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.
Specificity Specific for KCNS2.
Spcs X-react. Mouse, rat, human. Other species not yet tested.
Format Lyophilised
Reconstitution Reconstitute in 100 µl of sterile water. Centrifuge to remove any insoluble material.
Storage Maintain the lyophilised/reconstituted antibodies frozen at -20°C for long term storage and refrigerated at 2-8°C for a shorter term. When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles.
Expiry Date 12 months after reconstitution
Shipping This item will be shipped to you at ambient temperature in a lyophilised form.
Limitation For research use only



IHC-P on paraffin sections of rat brain stem.

The animal was perfused using Autoperfuser at a pressure of 130 mmHgwith 300 ml4% FA before beingprocessed for paraffin embedding. HIER: Tris-EDTA, pH 9 for20 minusing Thermo PT Module.

Blocking: 0.2% LFDM in TBST filtered thru 0.2 µm.

Detection was done using Novolink HRP polymer from Leica following manufacturers instructions; DAB chromogen: Candela DAB chromogen from Osenses.

Primary antibody: dilution 1: 3000, incubated 30 min at RT using Autostainer.

Sections were counterstained with Harris Hematoxylin.