

Rabbit antibody to TAS2R46

| ImmunogenA synthetic peptide from human TAS2R46 conjugated to blue carrier protein was used as the antigen.ConjugateUnconjugated antibodyAlso knownTaste receptor type 2 member 46, T2R46, T2R54, TAS2R54HostNZ white rabbitPurityWhole serumClonalityPolyclonalIsotypePolyclonal, whole serumApplicationsIHC, WB. A dilution of 1 : 1000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.SpecificitySpecific for TAS2R46.Spcs X-react.Human, mouse, rat, marmoset. Other species not yet tested.FormatLyophilisedReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and refrigerated at 2-8C for a shorter term. When reconstituting, glycerol (1:1) may be added for an |
|--|
| Also knownTaste receptor type 2 member 46, T2R46, T2R54, TAS2R54HostNZ white rabbitPurityWhole serumClonalityPolyclonalIsotypePolyclonal, whole serumApplicationsIHC, WB. A dilution of 1 : 1000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.SpecificitySpecific for TAS2R46.Spcs X-react.Human, mouse, rat, marmoset. Other species not yet tested.FormatLyophilisedReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| HostNZ white rabbitPurityWhole serumClonalityPolyclonalIsotypePolyclonal, whole serumApplicationsIHC, WB. A dilution of 1 : 1000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.SpecificitySpecific for TAS2R46.Spcs X-react.Human, mouse, rat, marmoset. Other species not yet tested.FormatLyophilisedReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| PurityWhole serumClonalityPolyclonalIsotypePolyclonal, whole serumApplicationsIHC, WB. A dilution of 1 : 1000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.SpecificitySpecific for TAS2R46.Spcs X-react.Human, mouse, rat, marmoset. Other species not yet tested.FormatLyophilisedReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| ClonalityPolyclonalIsotypePolyclonal, whole serumApplicationsIHC, WB. A dilution of 1 : 1000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.SpecificitySpecific for TAS2R46.Spcs X-react.Human, mouse, rat, marmoset. Other species not yet tested.FormatLyophilisedReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| IsotypePolyclonal, whole serumApplicationsIHC, WB. A dilution of 1 : 1000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.SpecificitySpecific for TAS2R46.Spcs X-react.Human, mouse, rat, marmoset. Other species not yet tested.FormatLyophilisedReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| ApplicationsIHC, WB. A dilution of 1 : 1000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.SpecificitySpecific for TAS2R46.Spcs X-react.Human, mouse, rat, marmoset. Other species not yet tested.FormatLyophilisedReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| end user. Not yet tested in other applications.SpecificitySpecific for TAS2R46.Spcs X-react.Human, mouse, rat, marmoset. Other species not yet tested.FormatLyophilisedReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| Spcs X-react.Human, mouse, rat, marmoset. Other species not yet tested.FormatLyophilisedReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| FormatLyophilisedReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| ReconstitutionReconstitute in 100 ul of sterile water. Centrifuge to remove any insoluble material.StorageMaintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| Storage Maintain the lyophilised/reconstituted antibodies frozen at -20C for long term storage and |
| |
| additional stability. Avoid freeze and thaw cycles. |
| Expiry Date 12 months after reconstitution |
| Shipping This item will be shipped to you at ambient temperature in a lyophilised form. |
| Limitation For research use only |